

Prevalence of Keratoconus and The Role of Advanced Optical Techniques for Correcting Vision

Dareen Alshareef Ahmed Jadullah¹, Aziza Al-Tarhouni², Fatima Walid Muhammad Al-Awami³, Mawda Tariq Mohammed Alfakhi³, Israa Salem Omar Al-Mshity³, Nour Al-Huda Hamad Muhammad Al-Zwai³

¹Lecturer at Pharmaceutical Technology Department, College of the Medical Technology, Benghazi-Libya. Email: dareen.shareef@cmtben.edu.ly

²Lecturer at Optometry & Vision Science department, College of the Medical Technology, Benghazi- Libya. Email: azizaatarhoni173@gmail.com

³Under Graduate Student at Department of Optometry & Vision Science, College of Medical Technology, Benghazi- Libya. Email: www.cmtben.edu.ly

الملخص: هدفت دراسة مقطعية، شملت مرضى مستشفى بنغازي التعليمي لطب العيون ومركز دلتا للبصريات، إلى تقييم مدى انتشار القرنية المخروطية (KC) ودراسة توزيع وفعالية التقنيات البصرية المتقدمة المستخدمة في علاجها. أجري استبيان شمل 500 مشارك، واستخدم برنامج SPSS لتحليل 413 استجابة صالحة. أظهرت النتائج أن القرنية المخروطية منتشرة بشكل كبير (59.6%)، وخاصة بين الفئات العمرية الأصغر. في حين لم تكن هناك علاقة ارتباطية ذات دلالة إحصائية بين القرنية المخروطية والعمر أو الجنس أو سُمك القرنية، فقد وُجدت علاقة ارتباطية إحصائية ذات دلالة إحصائية بين القرنية المخروطية والاستجماتيزم غير المنتظم ونوع العلاج. على الرغم من أن مرضى القرنية المخروطية كانوا أكثر عرضة للخضوع لإجراءات أكثر تعقيداً مثل تثبيت القرنية، وزرع حلقات داخل القرنية، وزراعة القرنية، إلا أن النظارات ظلت الوسيلة التصحيحية الأكثر شيوعاً. تُبرز هذه النتائج أهمية الكشف المبكر، وتنقيف المرضى، وتطبيق أحدث أساليب التشخيص والعلاج في رعاية العيون.

الكلمات المفتاحية: القرنية المخروطية؛ الانتشار؛ الربط المتقاطع للقرنية؛ الاستجماتيزم؛ المتقدم.

Abstract: A cross-sectional study, patients at the Benghazi Teaching Hospital for Ophthalmology and Delta Optics Center, aimed to assess the prevalence of keratoconus (KC) and investigate the distribution and efficacy of advanced optical techniques used to treat it. A survey with 500 participants was conducted, and SPSS was used to analyze 413 valid responses. The findings showed that keratoconus was quite prevalent (59.6%), specifically among younger age groups. While there was no significant correlation between KC and age, gender, or corneal thickness, there was a significant statistical correlation between KC and irregular astigmatism and the type of treatment. Although keratoconus patients were more likely to undergo more complex procedures like corneal cross-linking, intracorneal ring implantation, and transplantation, glasses remained the most common corrective measure. These findings highlight the importance of early screening, patient education, and the

application of state-of-the-art diagnostic and therapeutic approaches in ophthalmic care.

Keywords-; Keratoconus; prevalence; corneal crosslinking; astigmatism; advanced.

I. INTRODUCTION

1.1 Definition:

Keratoconus is a progressive, non-inflammatory eye disorder characterized by thinning and cone-like protrusion of the cornea, leading to irregular astigmatism and visual impairment. The condition typically begins in adolescence or early adulthood and can progress for 10 to 20 years before stabilizing. Fig (1). Its etiology is multifactorial, involving genetic, biochemical, and environmental factors, including chronic eye rubbing and atopy (Gomes *et al.*, 2015). Recent advancements in diagnostic technologies, such as corneal topography and tomography, have significantly improved early detection and classification of keratoconus. Furthermore, treatments like corneal collagen cross-linking (CXL) have proven effective in halting disease progression, reducing the need for corneal transplantation (Wollensak *et al.*, 2003; Hersh *et al.*, 2011).



Fig (1): Keratoconus of the Eye.

1.1 History Keratoconus:

Keratoconus was first reported by Benedict Duddell in 1736. (Wyman,1992); Following its first description,

various terminologies such as 2 relapses cornea, cornea conica, sugar- loaf cornea, and proclivaria cornea .(Grzybowski, 2013) Around a century later, John Nottingham furnished the primary designated description of the disorder in his landmark book in 1854 (Grzybowski, 2013; Gokul, 2016) Pickford defined the conical cornea as a sickness that is “intractable in nature and deadly to vision” and One in which” the pathology and remedy are so little understood. Around a hundred and seventy years later, keratoconus stays an enigmatic disease. Over the beyond few decades, speedy development in diagnosing and coping with keratoconus has been observed. Originally described as a tremendous illness with the resource of the use of the National Institute of Health with an incidence of plenty much less than 1 consistent with 2,000 People (Ferrari, 2020), it's far now acknowledged that keratoconus is a great deal greater not unusual place than firstly thought. The pronounced occurrence is pretty variable from 0.2 consistent with 100,000 in Russia (Gorshkova, 1998) to 33 consistent with 1,000 in Iran (Hashemi, 2013). A meta-analysis from 15countries reported a global prevalence of 1.4 per 1,000 (Hashemi, 2020). A better incidence is cited in Asian and Middle Eastern populations. Pediatric populations have a better occurrence fee, with a suggested occurrence fee starting from 5.2 consistent with 1,000 people In New Zealand to 47. nine consistent with 1,000 human beings in Saudi Arabia (Torres, 2018; Papali, 2019). In addition, it is one of the most now no longer unusual place signs for keratoplasty in plenty of countries. (Ting, 2012; Park, 2015) Nonetheless, a few nations have suggested a lowering fashion withinside the wide variety of keratoplasty for Keratoconus in view of the implementation of corneal cross- linking (Sandvik, 2015, Godefrooij, 2016).

1.2 Aim of This Study:

This study aimed to investigate the prevalence of keratoconus and assess the effectiveness and distribution of advanced optical techniques used in its treatment among patients attending the Benghazi Teaching Hospital for Ophthalmology and Delta Optics Center. Also, determine associations between keratoconus prevalence and demographic, anatomical, and treatment variables.

II. KERATOCONUS OVERVIEW AND LITERATURE REVIEWS

2.1. Genetics Causes:

Keratoconus has long been considered to have a genetic component, given its association with other genetic syndromes (such as Down's syndrome (Mathan, 2020) Leber's congenital amaurosis (Elder, 1994; Damji, 2001), Ehlers-Danlos syndrome (Robertson, 1975) and Noonan syndrome (Lee, 2014), its prevalence in first- degree relatives (Rabinowitz, 2021; Shaag A, 2013, Almusawi LA, 2021, Lapeyre G, 2020) and occurrence in monozygotic twins,(Edwards, 2001; Tuft SJ, 2012). It has been anticipated that a relative of an individual with keratoconus has a 15 to 67 times greater risk of developing keratoconus than an individual with no family history of keratoconus (Wang, 2000). Keratoconus follows an apparently autosomal dominant/recessive mode of inheritance in some families (Bisceglia, 2009; Gonzalez, 1996).

2.2. Cellular Biochemistry:

To date, 117 proteins and protein commands have been implicated withinside the pathophysiology of keratoconus. Differential expression of numerous corneal proteins effects in modifications withinside the structural integrity and morphology of the keratoconus cornea, through altering its collagen content and keratocyte apoptosis and necrosis in the (Yam, 2019; Srivastava, 2016). Oxidative stress markers and antioxidants are dysregulated in keratoconus, associated with an imbalance of redox homeostasis in tears, cornea, aqueous humour and blood (Navel, 2020). Keratoconus is related to a typical boom in oxidative strain markers, mainly in reactive oxygen and nitrogen species and malondialdehyde.

2.3. Biomechanical factors:

The degeneration of the proteoglycans across the stromal collagen fibrils in keratoconus corneas results in breakage of, and degeneration of the microfibrils within, collagen fibrils (Alkanaa, 2019). Erato-conic corneas have decreased levels of aldehyde dehydrogenase Class 3 (Gondhowiardjo, 1993) and superoxide dismutase enzymes (Behndig, 1998), Both enzymes play vital roles withinside the reactive oxygen techniques of various species. The reactive oxygen accumulation causes cytotoxic deposition of malondialdehyde and peroxynitrites, which could potentially damage corneal




Fig (3): Corneal halos and glare.

5. Photophobia:

A common symptom due to increased light scattering caused by the abnormal curvature of the cornea.

6. Poor Night Vision:

Many patients report difficulty seeing in the dark or in low-light environments, which is linked to higher-order optical aberrations.

7. Eye Irritation or Itching:

Patients may feel persistent irritation or itching, often associated with chronic eye rubbing — a major contributing factor. Within the improvement of the condition.

8. Monocular Diplopia or Multiple Images from One Eye:

This is a complex visual symptom where patients see double or multiple images from a single eye due to

abnormal corneal refraction, usually in superior levels of the disease sensation of Pressure or Mild Eye Discomfort. (Sykakis, 2015). In some cases, patients may feel a mild pressure or discomfort in the eye, particularly as the disease progresses. Sudden and Severe Vision Loss (in Case of Descemet's Membrane rupture).

2.7. Classification system of keratoconus:

Different class structures presently to be had for assessing keratoconus severity primarily based totally on:

- Corneal morphology and disorder evolution.
- Optical and visible function.
- Descriptors of corneal shape (i.e, index-primarily based totally systems).

2.7.1. Corneal morphology and disease evolution:

The maximum generally used type structures primarily based totally on morphological adjustments and ailment evolution are:

- Morphological (Buxton) classification— This device classifies the illness based completely on the shape and function of the cone into oval, nipple and globe keratoconus (Perry HD, 1980):

- (1) In oval keratoconus the cone impacts one or corneal quadrants.
- (2) Keratometric classification (Vega Estrada, 2017) – This system categorizes keratoconus into four grades based on the magnitude of the cornea's central corneal power: (a) Mild; (b) Moderate; (c) Advanced.
- (3) Hom's classification (Rabinowitz, 1980) – This system classifies keratoconus into four grades based on clinical signs: (a) Preclinical suggests that no keratoconus symptoms and symptoms are detected. (b) Mild cases display mild corneal thinning and scissors reflex. (c) Moderate suggests terrible visible great and corneal thinning without corneal scarring. Table (2).

Table (2): Amsler-Krumeich classification:

N o.	Amsler – Krumeich	Alio – Shabayek
1	Grade I Corneal steepening Refraction > 5 D Mean number one K readings < 48 D	No scar Coma-like RMS 1.50 to 2.50 μ m Mean number one K readings < 48 D
2	Grade II No scars Corneal thickness > four hundred μ m Refraction > eight D Mean valuable K readings < fifty-three D	No scars Corneal thickness > four hundred μ m Mean valuable K readings < fifty-three D Coma-like RMS > 2.50 to $\leq$ 3.50 μ m
3	Grade III No scars Corneal thickness > 300 μ m Refraction > 10 D Mean central K readings < fifty-five D	No scars Corneal thickness > 300 μ m Coma-like RMS > 3.50 to $\leq$ 4.50 μ m Mean central K readings < fifty-five D
4	Grade IV Central scarring Corneal thickness > 200 μ m Not reliable refraction Mean central K readings > fifty-five D	Central scarring Corneal thickness > 200 μ m Coma-like RMS > 4.50 μ m Mean central K readings > fifty-five D

2.7.2. Optical and visual function:

This classification associated with a significant decrease in optical quality resulting from increases in ocular aberrations and a loss of corneal transparency in some cases which can affect quality of life (Kandel, 2020).

- Lio-Shabayek (2006) - This system, which is based on the Amsler–Krumeich classification.
- Keratoconus Severity Score (KSS) (McMahon, 2006) - This system grades the severity of keratoconus from 0 (suspect) to 5 (severe) primarily based totally on corneal topographic indices (i.e., anterior corneal better order aberration RMS blunders and suggest relevant keratometry).
- RETICS classification (Ali'o, 2011) – In addition to clinical signs and optical and visual function variables, this category device additionally takes into attention corneal biomechanical parameters (i.e., hysteresis and resistance factor).
Belin ABCD grading system (Belin, 2016) – Keratoconus severity is graded primarily based totally on 4 variables:
 - Anterior and posterior corneal radius.
 - Curvature of the 3.0 mm central zone of the thinnest corneal location.
 - Thinnest pachymetry.
 - Distance best corrected visual acuity.

2.7.3. Index-based systems:

These structures might also additionally consist of one or extra variables for keratoconus detection and commonly use cut-off values to permit differentiation among regular corneas, keratoconus suspects, and clinical keratoconus four hundred μm .

2.8. Stages of Keratoconus:

Keratoconus progresses through a series of clinical stages, each characterized by increasing corneal deformation and visual impairment. The classification into stages helps guide treatment and monitor disease progression. The most widely accepted staging system is based on corneal curvature, pachymetry (thickness), visual acuity, and topographic findings.

Stage 1 –Early stage:

- Slight corneal thinning, mostly inferiorly.
- Irregular astigmatism begins to appear.
-

- Best-corrected visual acuity (BCVA).
- Still good with glasses.
- Topography shows mild asymmetry or inferior steepening.

Stage 2 – The Moderate Stage:

- The corneal protrusion and thinning increased.
- Visual acuity worsens; glasses are no longer sufficient, and rigid gas permeable lenses may be required.
- Keratometric readings (K-values) between 45–52 diopters.
- Topographic signs include a pronounced cone shape and possible scissoring reflex on retinoscopy.

Stage 3 – The Advanced Stage:

- Corneal thinning and ectasia have been noted.
- Vision is significantly reduced; contact lenses may no longer provide stable correction.
- K-readings typically between 52–62 diopters.
- Corneal haze may begin to appear.

Stage 4 – The End-stage:

- Thinning and scarring of the cornea noted to be aggressive.
- Vision is severely affected and often uncorrectable with lenses.
- K-readings exceed 62 diopters.
- Corneal transplantation as the surgical treatment may use at this stage.

2.9. Diagnosis:

2.9.1. Examination:

Scissoring of the red reflex on retinoscopy is a dependable and touchy technique for detecting early-degree KC. External signs consist of the Munson sign (V-formed deformation of the decrease eyelid because of the cone whilst the affected person appears down; And Rizzuti sign (conical illumination at the nasal sclera whilst light is directed at the cornea from the temporal side..(However, these external signs are typically not observed in mild KC (Hashemi H, 2020).

2.9.2. Topography and tomography:

2.9.2.1. Corneal Topography:

Topography permits noninvasive qualitative and quantitative characterization of corneal morphology.

Topographic maps will display abnormal astigmatism with steepening. The following maps are analyzed: anterior, sagittal, and tangential curvature maps; anterior and posterior elevation maps; and the thickness map (Martinez-Abad, 2017).

- **Pentacam Topography:**

Is a diagnostic imaging technique used in ophthalmology to analyze the cornea and anterior segment of the eye. It utilizes a Scheimpflug camera to create detailed maps of the cornea's surface, including its curvature, elevation, and thickness. Figure (4).

- **Autorefractometer:**

Refractometry or optometry is topographic assessing refractive error with a refractometer or optometer instrument. Automated refractors or autorefractors are instruments designed Figure (5), to assess the refractive error and can vary based on the underlying principle (Xiong S, 2017).

2.9.2.2. Corneal Tomography:

Tomography provides additional parameters for evaluating the anterior and posterior corneal surfaces. Early posterior corneal structural changes, including stromal thinning and elevation changes, are observed prior to anterior surface changes in kc.4 this allows for reliable detection of early-stage kc even before a patient becomes symptomatic.



Fig (4): Pentacam Devices.



Fig (5): Autorefractometer Devices.

2.9.3. Other Techniques:

2.9.3.1. Ocular Response Analyzer (Reichert):

Evaluates corneal biomechanics by measuring corneal hysteresis, the difference in applanation pressure when the cornea bends inward in response to a jet of air and when it returns to its normal state (Martinez-Abad, 2017).

2.9.3.2. I Trace aberrometer:

I TRACE is a ray tracing aberrometer which combines both the wavefront aberrometer as well as Placido based corneal topography (Sinha A, 2019). Figure (6)



Fig (6): I Trace aberrometer.

2.10. Management and Treatment:

2.10.1. Mild keratoconus:

Spectacles can only be used in mild cases of keratoconus, and often result in poor visual acuity (Rabinowitz YS, 1998). Figure (7).

2.10.2. Moderate keratoconus:

Currently, it is estimated that 90% of patients affected by corneal irregularity utilize contact lenses (Zadnik K, 1998).




Fig (7): Eye Glasses Keratoconus.




Fig (8): Scleral contact lenses.

2.10.2.1. Rigid Corneal Contact Lenses:

Rigid corneal touch lenses and piggyback systems. Three techniques had been historically used for becoming inflexible corneal touch lenses in keratoconus which includes apical clearance, apical touch, and three-point touch (Romero-Jiménez, 2013; Romero-Jiménez M, 2015).

2.10.2.2. Corneoscleral and Scleral Lenses:

Corneoscleral lenses are described as any inflexible touch lens with shared bearing among the peripheral cornea and conjunctiva overlying the sclera, irrespective of the overall lens diameter (De Luis, 2018). The principal benefits of this lens designs in comparison with inflexible corneal lenses are stepped forward consolation because of the decreased lens edge-eyelid interaction and enhanced stability and centration with larger optical zones for more consistent vision across a range of pupil diameters (Downie, 2015). Figure (8).

2.10.2.3. Soft Contact Lenses:

Soft lenses are to be had in excessive round and toric powers for the correction of myopia and astigmatism in early keratoconus, decentered cones, and for patients with rigid lens intolerance (Yilmaz, 2016; Sultan, 2016).

2.10.2.4. A Hybrid Contact Lens:

A hybrid contact lens consists of a rigid corneal lens and a peripheral soft skirt to combine the optical benefits of corneal rigid lenses and the comfort provided by soft contact lenses. Early generation hybrid lenses were often associated with decreased comfort, complications due to the use of low oxygen permeability materials, and reduced durability of the GP/soft material interface (Rubinstein MP, 1991; Leal F, 2007; Pilskalns 2012).

2.10.3. Severe keratoconus:

Severe instances of keratoconus can be controlled with scleral lenses, mainly while different lens modalities normally fail to achieve a physiologically acceptable fit (Ling JJ, 2021; Koppen C, 2018; Segal O, 2003; Rosenthal P, 2005).

2.10.3.1. Surgical Procedures:

2.10.3.1.1. Cross-linking Corneal (CXL):

Cross-linking increases the (Santodomingo-Rubido *et al.*, 2017), biomechanical stability and rigidity of the cornea in an attempt to prevent keratoconus progression. The method includes the elimination of imperative 6–7 mm of corneal epithelium accompanied via way of means of the following software of 0.1% riboflavin solution and corneal radiation of ultraviolet-A mild at 370 nm (Sorkin, 2014; O'Brart, 2017; Beckman, 2019). Figure (9).



Fig (9): Corneal Cross Linking.

2.10.3.1.2 Refractive Surgery:

Various refractive surgical operation interventions had been used for keratoconus management, with phakic lens implantation and photorefractive keratectomy (PRK), being the two most widely studied (Kim KH, 2020; Fern´andez-Vega-Cueto L, 2017)

1. Photorefractive keratectomy of the Corneal (PRK):

It makes use of an excimer laser to completely regulate the form of the anterior vital cornea via way of means of getting rid of a small segment of stromal tissue via way of means of vaporization. outcomes in keratoconic eyes were fairly successful, with a few research staring at a vast discount in cone evolution in incipient cases (Kasparova, 2002).

2. Toric intraocular lens implantation (IOL):

Phakic and pseudophakic intraocular lens implantation for the remedy of keratoconus is typically executed together with different corneal refractive surgical treatment methods, such as corneal rings or keratoplasty (Peˆna-García, 2015).

2.10.3.1.3. Corneal Transplantation and Implantation:

Corneal transplantation is the conventional remedy for superior keratoconus has been suggested to be the cause for 18% of penetrating keratoplasty procedures, and 40% of deep anterior lamellar keratoplasty interventions (Gadhvi , 2019; Amalich-Montiel, 2016). Figure (10).



Fig (10): Corneal Transplantation.

1. Keratoplasty:

Penetrating keratoplasty (PK), which includes the elimination of the whole thickness of the cornea and alternative with donor tissue (Brierly SC, 2000). Deep

anterior lamellar keratoplasty (DALK) is every other surgical method used to update diseased recipient stroma with donor corneal stroma, whilst the recipient corneal endothelium and posterior restricting lamina are retained.

2. Anterior proscribing lamina transplantation:

Anterior limiting lamina transplantation is a novel technique that may stabilize progressive ectatic corneal changes in eyes with advanced keratoconus, which are too steep or too thin for CXL or ICRS (Brierly SC, 2002).

3. Implantation of stem cells (Intrastromal):

There is Two methods of intrastromal implantation, first implanting of stem cells with a biodegradable scaffold; second of stem cells with a non- biodegradable scaffold; and intrastromal implantation of stem cells with a decellularized corneal stromal scaffold (Amalich- Montiel, 2016; Ali´o, 2020).

• Ring Segments (Intracorneal):

Intracorneal ring segments (ICRS) are small PMMA (polymethyl methacrylate) devices, which might be implanted into the cornea aiming to alter its geometry and enhance its refractive homes and patient’s visible acuity. Colin delivered using ICRS implantation for the control of keratoconus in 2000 (Colin J,2000). Nevertheless, Reynolds become the primary to implant a 360° intracorneal ring for the control of myopia in 1978 (Burris,1998). Figure (11).



Fig (11): Intracorneal Ring.

2.11. Literature Reviews:

In Kenya a prospective, a cross-sectional study Aly Rashid Z, H *et al* about the Prevalence and demographic profile of keratoconus among high school students in Kenya, from 25th October 2021 to the 26th August 2022 through Using a stratified cluster random sampling

method. As students diagnosed with visual acuity measurement, auto-refraction, retinoscopy and corneal

topography. (Rashid *et al.*, 2021). Their result shows 57.3% where female, also had disease prevalence of KC and KCS to be 1.7% and 3.8% respectively, also with no significant relationship between prevalence of keratoconus and (gender and age) (Aly Rashid *et al.*, 2025).

In study by Kenyan authors Aly Rashid and colleagues were used an online questionnaire distributed among ophthalmic clinical officers and optometrists, from 13th September to 28th October 2021, used retinoscopes, slit lamps, corneal topographer as diagnosed methods in their study, as majority of respondents 90.2% in their study treated with spectacles in mild cases, 29% in moderate cases and only 1.9% in severe cases and 26.1% of patients with keratoconus referred for CXL (Aly Rashid *et al.*, 2023).

A cross-sectional Sweden study by Binder J.T, Vibeke Z.H aimed to assess prevalence and incidence of keratoconus from 1st January, 2010, to 31st December, 2020, the prevalence of keratoconus in their study was 169.5 per 100,000, however 75% of them were male. Also, the prevalence was highest in the age group 21 to 30 years, 348.4 per 100,000 (Binder, 2023). However, a Danish study by Bak-Nielsen *et al.*, had prevalence of keratoconus was estimated to be 3.6 per 100,000 annually. Additionally, most of keratoconus Patients were male (Bak- Nielsen *et al.*, 2025).

III. METHODOLOGY

3.1. Study Design and Population:

- Its cross-sectional descriptive study was conducted aimed to explore the prevalence of keratoconus and assess the effectiveness and distribution of advanced optical techniques used in its treatment among patients attending the Benghazi Teaching Hospital for Ophthalmology and Delta Optics Center, from 1st of May to 26th of June.
- A total of (500) patients enrolled in this study from both genders; the process of patient's selection was randomly according to their presence at hospital or center. A structured questionnaire was administered to (500) individuals, out of which (413) valid responses were obtained and included in the final analysis and seven patients were excluded.

3.2. Data Collection Tool:

The process of data collection was through questionnaire self- made based on previous studies. The questionnaire collected: Demographic information (age and gender), clinical characteristics (corneal thickness, type of astigmatism), and details about keratoconus diagnoses through clinical examination or pentacam and methods of treatment used (eyeglasses, contact lenses, cross-linking, corneal transplant, or ring implantation).

3.3. Data Analysis:

Data analysis was performed by using the Statistical Package for the Social Sciences (SPSS). Descriptive data described as Frequencies and percentages. The Pearson Chi-square test was used to determine the relationships between the prevalence of keratoconus and other variables in the study. A significance level of 0.05 was used for all statistical tests.

3.4. Ethical Approval:

The study was approved from Medical Technology college - Optometry & Vision Science department, all the research procedures were carried out for patients after consent with ethical guidelines. Also was permitted from Benghazi Teaching Hospital for Ophthalmology, and Delta Optics Center.

IV. RESULTS

4.1. Descriptive Results:

In an attempt to reinforce what was discussed in the theoretical aspect of this project, a questionnaire was prepared and distributed to 500 people in both Benghazi Teaching Hospital for Ophthalmology, and Delta Optics Center.

413 questionnaires were received, all of which were valid. This data was analyzed using the statistical analysis program SPSS and a chi-square test was conducted to try to describe the relationships between the different variables.

Table (3), and Figure (11): Illustrates the respondents' distribution according to the keratoconus prevalence. It reveals that there were 246 prevalent patients, or 59.6% of the total, and 167 non-prevalent patients, or 40.4%.

Table (3): The distribution of respondent's base on the prevalence of KC:

Keratoconus	umber of respondents	%
Prevalent	246	59.6
Non- prevalent	167	40.4
Total	413	100

The distribution of respondents based on the prevalence of Keratoconus



Fig (11): The distribution of respondents based on the prevalence of KC.

Table (4), and Figure (12): Shows the distribution of respondents based on the Gender. It shows that the number of Male patients had greater Keratoconus prevalence 232 (56.2%) than female 181(43.8%).

Table (4): The distribution of respondent's base on the Gender:

Gender	Number of respondents	%
Male	232	56.2
Female	181	43.8
Total	413	100

The distribution of respondents based on the Gender



Fig (12): The distribution of respondents based on the Gender.

Table (5), and Figure (13): It is evident that the age group of [15–24] had the largest percentage of respondents (44.6%), followed by [25–34] (21.8%), [45–54] (18.2%), and [35–44] (15.5%) respectively.

Table (5): The distribution of respondents based on the Age Interval:

Age interval	Number of respondents	%
15-24	184	44.6
25-34	90	21.8
35-44	64	15.5
45-54	75	18.2
Total	413	100

The distribution of respondents based on the age intervals




Fig (13): The distribution of respondents based on the age intervals.

Table (6), and Figure (14): Display the respondents' distribution by corneal thickness, with the largest percentage of respondents (71.9%) falling into the "More than 500" category and the lowest number (28.1%) falling into the "Less than 500" category.

Table (6): The distribution of respondents based on the Corneal Thickness:

Corneal Thickness	Number of respondents	%
More than 500	297	71.9
Less than 500	116	28.1
Total	413	100

The distribution of respondents based on the Corneal Thickness




Fig (14): The distribution of respondents based on the Corneal Thickness.

Table (7), and Figure (15): It clear that, the percentage of 59.1% falls in category of irregular Astigmatism, while the percentage of 40.9% falls in category of regular Astigmatism.

Table (7): The distribution of respondents based on the Astigmatism:

Astigmatism	Number of respondents	%
Regular	169	40.9
Irregular	244	59.1
Total	413	100

The distribution of respondents based on the Astigmatism




Fig (15): The distribution of respondents based on the Astigmatism.

Table (8), and Figure (16): Demonstrating the respondents' distribution by treatment, it is evident that the eyeglasses category accounted for 43.8% of the total, followed by the corneal cross-linking category with 21.8%, contact lenses with 16.5%, intra-corneal ring implantation with 9.2%, and corneal transplantation with 8.7%.

Table (8): The distribution of respondents based on the Treatment.

Treatment	Respondents	%
Eyeglasses	181	43.8
Contact Lenses	68	16.5
Corneal Cross Linking	90	21.8
Corneal Transplantation	36	8.7
ICR Implantation	38	9.2
Total	413	100

The distribution of respondents based on the Treatment




Fig (16): The distribution bas respondents based on the Treatment.

Table (9), and Figure (17): Represent the distribution of the respondents based on the prevalence of Keratoconus and their gender, it seems that, the number of Male prevalence is 131, and the number of Female prevalence is 115.

Table (9): The distribution of respondents based on the prevalence of Keratoconus and their gender:

Keratoconus	Gender		Total
	Male	Female	
Prevalence	131	115	246
Non-prevalence	101	66	167
Total	232	181	413

The distribution of respondents based on the prevalence of Keratoconus and their gender




Fig (11): the distribution of respondents based on the prevalence of Keratoconus and their gender.

4.2. Statistical Result:

Table (10): It seems that; at ($\alpha=0.05$) there is no statistically significant differences between the gender of the respondent and the prevalence of Keratoconus, whereas (Chi-square = 2.110), and P-value (0.146) > 0.05.

Table (10): Chi-square test Gender and prevalence of KC:

	Value	d.f.	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.110	1	0.146

Table (11), and Figure (18): Show that the number of Prevalence respondents of age interval [15-24] was 110, then 49 for age interval [25-34], followed by 42 for age interval [35-44], and lastly 45 for age interval [45-54]. While the number of non-prevalence keratoconus respondents of age interval [15-24] was 74, followed 41 for age interval [25-34], then 22 for age interval [35-44], and lastly 30 for age interval [45-54].

Table (11): The distribution of respondents based on the prevalence of KC, and their age interval:

Keratoconus	Age Interval				Total
	15-24	25-34	35-44	45-54	
Prevalence	110	49	42	45	246
Non-Prevalence	74	41	22	30	167
Total	184	90	64	75	413

The distribution of respondents based on the prevalence of Keratoconus, and their age interval




Fig (18): The distribution of respondents based on the prevalence of Keratoconus, and their age interval.

Table (12): It seems that; at ($\alpha=0.05$) there is no statistically significant differences of the relationship between the age interval of the respondent and the prevalence of Keratoconus, whereas (Chi-square = 1.965), and P-value (0.580) > 0.05.

Table (12): Chi-square test:

	Value	d.f.	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.965	3	0.580

Table (13): The distribution of the Prevalence of Keratoconus, and the Corneal Thickness:

Corneal Thickness		Keratoconus	Total
> 500	Prevalence	76	246
< 500	Non-Prevalence	40	167
	Total	116	413

The distribution of the Prevalence of Keratoconus, and the Corneal Thickness




Fig (19): The distribution of the Prevalence of Keratoconus, and the Corneal Thickness.

Table (13), and Figure (19): Display the amount of prevalence of Keratoconus with > 500 Corneal Thickness is 170, and the amount of prevalence of Keratoconus with < 500 Corneal Thickness is 76. However, the number of non-prevalence of Keratoconus with > 500 Corneal Thickness is 127, and the number of non-prevalence of Keratoconus with < 500 Corneal Thickness is 40.

Table (14): It seems that; at ($\alpha=0.05$) there is no statistically significant differences of the relationship between the Corneal Thickness of the respondent and the prevalence of Keratoconus, whereas (Chi-square = 2.373), and P-value (0.123) > 0.05.

Table (15), and Figure (20): Display the amount of prevalence of Keratoconus and Regular Astigmatism is 10, and the amount of prevalence of Keratoconus and Irregular Astigmatism is 236. However, the number of non-prevalence of Keratoconus regular Astigmatism is 159, and the number of non-prevalence of Keratoconus and Irregular Astigmatism is 8.

Table (14): Chi-square test:

	Value	d.f.	Asymp. Sig.
Pearson Chi-Square	2.373	1	0.123

Table (16): Chi-square test.

	Value	d.f.	Asymp. Sig. (2-sided)
Pearson Chi-Square	341.811	1	0.000

Table (15): The distribution of the Prevalence of Keratoconus, and the Astigmatism:

Keratoconus	Astigmatism		Total
	Regular	Irregular	
Prevalence	10	236	246
Non-Prevalence	159	8	167
Total	169	244	413

The distribution of the Prevalence of Keratoconus, and the Astigmatism




Fig (20): The distribution of the Prevalence of Keratoconus, and the Astigmatism.

Table (15): It seems that; at ($\alpha=0.05$) there are statistically significant differences of the relationship between the Astigmatism of the respondent and the prevalence of Keratoconus, whereas (Chi-square = 341.811), and P-value (0.000) < 0.05 .

Table (16), and Figure (21): Display the amount of prevalence of keratoconus and eyeglasses treatment, which was 65, followed by the amount of prevalence of keratoconus and contact lenses treatment, which was 60; then the amount of prevalence of keratoconus and corneal cross-linking treatment 63; then the amount of prevalence of keratoconus and corneal transplantation 30; and lastly the amount of prevalence of intra- corneal ring implantation treatment was 28.

According to Table (17), the number of non-prevalence of keratoconus and eyeglasses treatment was 116, followed by the number of non-prevalence of keratoconus and contact lens treatment, which was 8, then the number of non-prevalence of keratoconus and corneal cross-linking treatment was 27; then number of non-prevalence of keratoconus and corneal transplantation was 6; and lastly, the number of non-prevalence of keratoconus and intra-corneal ring implantation treatment was 10.

The distribution of respondents based on the prevalence of Keratoconus, and Treatment




Fig (21): The distribution of respondents based on the prevalence of Keratoconus, and Treatment.

Table (17): The distribution of respondents based on the prevalence of Keratoconus, and Treatment:

Keratoconus	Treatment					Total
	Eyeglasses	Contact Lenses	Corneal Cross Linking	Corneal Transplantation	Intra-corneal Ring Implantation	
Prevalence	65	60	63	30	28	246
Non-Prevalence	116	8	27	6	10	167
Total	181	68	90	36	38	413

Table (18): It seems that; at ($\alpha=0.05$) there are statistically significant differences of the relationship between the Treatment of the respondent and the prevalence of Keratoconus, whereas (Chi-square = 80.910), and P-value ($0.000 < 0.05$).

Table (18): Chi-square test:

	Value	d.f.	Sig. (2- sided)
Pearson Chi-Square	80.910	4	0.000

V. DISSCUSSIONS

“Keratoconus has been found to affect all ethnicities, although the prevalence and incidence are higher among South Asians and Middle Easterners compared with those of European ancestry.” (Kanski, 2015). The intent of this study was to investigate the probability of keratoconus among patients attend visit Delta Optics Center and Benghazi Teaching Hospital for Ophthalmology, as well as any potential correlations between the condition and clinical and demographic factors.

The result shows that keratoconus affected about 59.6% of patients, indicating a relatively significant disease burden in this area. Despite statistical analysis using the chi-square test showing no significant association between gender and keratoconus prevalence ($p = 0.146$), males had a slightly higher prevalence of keratoconus (53.3%)

than females. In contrast higher prevalence of KC in this study compared to study reported in kenya, Denmark.

Also, a Saudi Arabia study demonstrate that majority of middle east countries had higher

prevalence KC (Kennedy, 2000; Assiri, 2005). Opposite to this a study by Aly Rashid *et al.*, 2021) Kenyan study where female had higher prevalence of keratoconus over male, while some studies shown no difference between gender (Gomes, 2022), in compatible with this study by Aly Rashid Z, H et al. where prevalence of KC was not significantly associated with gender ($p = 0.80$). (Aly Rashid, 2023).

However, estimated prevalence of KC to be 7.9% in Africa a meta-analysis of a finite number of studies by Akowuah (Akowuah *et al.*, 2021), as consider lower when compared to this study.

Also, a lower prevalence of KC ranges from 1.4 to 4.8% In Turkey, India and the Middle East (Salman A, 2022; Özalp O, 2021; Jonas, 2009; Torres Netto, 2018). Concerning the age of participants as found, the 15–24 age group had the highest prevalence (44.7%), which is consistent with research demonstrating keratoconus often begins in adolescence or early adulthood. Even if younger people are more impacted, age by itself may not be a predictive element in this group also, relationship between 34 Statistically, age groups and the prevalence of keratoconus were not significant ($p = 0.580$). This suggests that other variables like heredity, environmental exposures, or eye habits may be more important predicting elements for keratoconus than age alone.

Similarly, a study by Kenyan authors Aly Rashid and colleagues found non-significant association between frequency of keratoconus and age (Aly Rashid *et al.*, 2025). By contrast in a cross-sectional study in Sweden, where prevalence was higher in the age group of 21 to 30 years (Binder, 2020), implying that regional

variance may be explained by variations in population characteristics, environmental conditions, and genetic predisposition. Age and gender alone cannot be regarded a deciding factor. The variances seen between several studies draw attention to the impact of genetic and environmental variation between populations.

In this study about 71.9% of the participants had a thickness greater than 500 microns, also, however still the correlation between corneal thickness and keratoconus prevalence found not significant ($p = 0.123$). This might indicate that although corneal thinning is a characteristic of keratoconus, it is not the only factor that determines early diagnosis or detection in this disease.

A statistically significant association was remarkably noted between keratoconus and astigmatism ($p < 0.001$). In 95.9% of patients with keratoconus had irregular astigmatism, which is consistent with the disease's known clinical profile. This result demonstrated how crucial comprehensive screening for astigmatism is in detecting possible cases of keratoconus as discussed before in theoretical part.

Eventually, as marked in the results a significant relationship was found between the type of treatment used and keratoconus prevalence ($p < 0.001$). Contact lenses, corneal cross-linking, and surgical procedures like corneal transplantation and intracorneal ring implantation were more common among keratoconus patients, even though eyeglasses were still the most popular treatment for all groups.

Also, in Sweden Kenyan study had found majority of the respondents prescribed spectacles in mild cases of keratoconus fewer patients where on for contact lenses and CXL and surgery treatment (Aly Rashid ,2025; Aly Rashid, 2023; Binder J.T,2020). Overall, the relatively higher prevalence incident reflects "the prevalence of KC appears 35 to be higher in countries with hot and dry climates compared to cooler climates" (Santodomingo-Rubido, 2022).

This distribution reflects both the severity and progression of the disease in those affected. This finding only represents patients in Benghazi and is not entirely conclusive because it only applies to

one hospital and clinic and does not represent the entire Libyan population.

CONCLUSION

Keratoconus is relatively common among patients visiting ophthalmology centers in Benghazi, particularly among young people. While gender, age, and corneal thickness were not significantly associated with the disease, irregular astigmatism had a strong and statistically significant relationship. Furthermore, the treatment options for keratoconus and non-keratoconus patients differed significantly, with the former receiving more advanced interventions.

This result represents the population in Benghazi not the entire country of Libya. Additionally, furthermore studies required in despite sample size, because the majority of participants come from a single hospital and clinic, cause sampling locations are not diverse. As noted, this study emphasizes the need for early screening for astigmatism in young people to help with early diagnosis and treatment. The study also supports the idea that treatment should be different for each patient based on disease degree and condition.

RECOMMENDATIONS

1. Encourage younger adults for early screenings regarding keratoconus specially for patients with irregular astigmatism, for early diagnosis and treatment, and emphasize them for continuous follow-up as irregular astigmatism consider related to keratoconus.
2. Increase population awareness through learning courses and programs regarding keratoconus signs and symptoms, risk factor so as to promote early diagnosis and treatment.
3. Improve the education of the eye care providers through training learning sessions about modern optical techniques and how to keep up with modernity about keratoconus.
4. Emphasize the important of conduct continuous studies related to keratoconus in Libya, address the updated concerning the risk factors, causes of disease.
5. Offer in public health care institutions updated diagnostic instruments and treatment options.

ACKNOWLEDGMENT

We thank Allah for His blessings. We sincerely appreciate our supervisor, Dr. Dareen Al-Sharif Ahmed Jadallah, for her guidance and support. We also thank Dr. Abdul Fattah Al-Mahdawi, Ms. Zainab Al-Haiblo, and the staff of Al-Sabri Polyclinic and Al-Nahr Central Hospital for their cooperation.

REFERENCES

- Ahuja P, Dadachanji Z, Shetty R, Nagarajan SA, Khamar P, Sethu S, *et al.* Relevance of IgE, allergy and eye rubbing in the pathogenesis and management of Keratoconus. *Indian J Ophthalmol* 2020; 68:2067–74.
- Akowuah PK, Kobia-Acquah E, Donkor R, Adjei-Anang J, Ankamah-Lomotey S (2021) Keratoconus in Africa: a systematic review and meta-analysis. *Ophthalm Physiol Opt* 41(4):736–747.
- Ali'ó del Barrio JL, Arnalich-Montiel F, De Miguel MP, El Zarif M, Ali'ó JL. Corneal stroma regeneration: Preclinical studies. *Exp Eye Res* 2021.
- Ali'ó JL, Piñero DP, Ales'ón A, Teus MA, Barraquer RI, Murta J, *et al.* Keratoconus-integrated characterization considering anterior corneal aberrations, internal astigmatism, and corneal biomechanics. *J Cataract Refract Surg* 2011; 37:552–68.
- Ali'ó JL, Shabayek MH. Corneal higher order aberrations: A method to grade keratoconus. *J Refract Surg* 2006; 22:539–45.
- Alkanaa A, Barsotti R, Kirat O, Khan A, Almubrad T, Akhtar S. Collagen fibrils and proteoglycans of peripheral and central stroma of the keratoconus cornea– ultrastructure and 3D transmission electron tomography. *Sci Rep* 2019; 9:19963.
- Almusawi LA, Hamied FM. Risk factors for development of keratoconus: a matched pair case-control study. *Clin Ophthalmol* 2021; 15:3473–9.
- Aly Rashid Z.H *et al.* Diagnosis and management of keratoconus by eye care practitioners in Kenya. *BMC Ophthalmology* (2023) 23:37.
- Aly Rashid Z.H *et al.* Prevalence and demographic profile of keratoconus among high school students in Kenya. *Int Ophthalmol.*(2025)
- Amsler M. K'eratocône classique et k'eratocône fruste; arguments unitaires. *Ann Ocul (Paris)* 1947; 180:112. 39.
- Amsler, M. (1950). *Le k'eratocône fruste au Javal. Revue d'Optique Théorique et Instrumentale*, 29, 721–736. (The earliest classification system for subclinical keratoconus).
- Arnalich-Montiel F, Ali'ó del Barrio JL, Ali'ó JL. Corneal surgery in keratoconus: which type, which technique, which outcomes? *Eye Vis* 2016; 3:2. doi: 10.1186/s40662-016-0033-y.
- Assiri AA, Yousuf BI, Quantock AJ, Murphy PJ: Incidence and severity of keratoconus in Asir province, Saudi Arabia. *Br J Ophthalmol.* 2005, 89:1403-6. 10.1136/bjo.2005.074955
- Bak-Nielsen, S., Ramlau-Hansen, C. H., Ivarsen, A., Plana-Ripoll, O., & Hjortdal, J. (2019). nationwide population-based study of social demographic factors, associated diseases and mortality of keratoconus patients in Denmark from 1977 to 2015. *Acta Ophthalmologica*, 97. Jonathan Binder,1,2 Vibeke Sundling.
- Beckman KA, Gupta PK, Farid M, Berdahl JP, Yeu E, Ayres B, *et al.* Corneal crosslinking: Current protocols and clinical approach. *J Cataract Refract Surg* 2019; 45:1670–9.
- Behndig A, Svensson B, Marklund SL, Karlsson K. Superoxide dismutase isoenzymes in the human eye. *Invest Ophthalmol Vis Sci* 1998; 39:471–5.
- Belin MW, Duncan JK. Keratoconus: The ABCD grading system. *Klin Monbl Augenheilkd* 2016; 233:701–7.
- Belin, M. W., & Duncan, J. K. (2016). Keratoconus: *The ABCD grading system. Keratoconus Symposium Abstracts*, ASCRS Annual Meeting.
- Binder J.T, Vibeke Z.H Prevalence and incidence of keratoconus in Sweden: A nationwide register

study between 2010 and 2020. *SJOVS*, June 2023.

Bisceglia L, De Bonis P, Pizzicoli C, Fischetti L, Laborante A, Di Perna M, *et al.* Linkage analysis in keratoconus: Replication of locus 5q21.2 and identification of other suggestive loci. *Investig Ophthalmol Vis Sci* 2009; 50:1081–6.

Brierly SC, Izquierdo L, Mannis MJ. Penetrating keratoplasty for keratoconus. *Cornea* 2000; 19:329–32.

Burris TE. Intrastromal corneal ring technology results and indications. *Curr Opin Ophthalmol*. 1998; 9:9–14.

Bykhovskaya Y, Rabinowitz YS. Update on the genetics of keratoconus. *Exp Eye Res* 2021; 202:108398.

Colin J, Cochener B, Savary G, *et al.* Correcting keratoconus with intracorneal rings. *J Cataract Refract Surg*. 2000; 26:1117–22.

Damji KF, Sohocki MM, Khan R, Gupta SK, Rahim M, Loyer M, *et al.* Leber's congenital amaurosis with anterior keratoconus in Pakistani families is caused by the Trp278X mutation in the AIPL1 gene on 17p.

Davidson, A. E., Hayes, S., Hardcastle, A. J., & Tuft, S. J. (2014). The pathogenesis of keratoconus. *Eye*, 28(2), 18G–1G5.

De Luis EB, Etxebarria Ecenarro J, Santamaria Carro A, Feijoo LR. Irregular corneas: Improve visual function with scleral contact lenses. *Eye Contact Lens* 2018; 44:159–63.

Downie LE, Lindsay RG. Contact lens management of keratoconus. *Clin Exp Optom* 2015; 98:299–311.

Edwards M, McGhee CNJ, Dean S. The genetics of keratoconus. *Clin Exp Ophthalmol* 2001; 29:345–51.

Elder MJ. Leber congenital amaurosis and its association with keratoconus and keratoglobus. *J Pediatr Ophthalmol Strabismus* 1994; 31:38–40.

Fernández-Vega-Cueto L, Romano V, Zaldivar R, Gordillo CH, Aiello F, Madrid-

Costa D, *et al.* Surgical options for the refractive correction of keratoconus: myth or reality. *J Ophthalmol* 2017; 2017:7589816.

Ferrari G, Rama P, The keratoconus enigma: A review with emphasis on pathogenesis, *The Ocular Surface* 2020, 18:363-373.

Ferrari G, Rama P. The keratoconus enigma: a review with emphasis on pathogenesis. *Ocul Surf.* (2020) 18:363–73. doi: 10.1016/j.jtos.2020.03.006.

Göncü T, Akal A, Adibelli FM, Çakmak S, Sezen H, Yilmaz ÖF. Tear film and serum prolidase activity and oxidative stress in patients with keratoconus. *Cornea* 2015; 34:101–23.

Gadhvi KA, Romano V, Fernández-Vega Cueto L, Aiello F, Day AC, Allan BD. Deep anterior lamellar keratoplasty for keratoconus: Multisurgeon results. *Am J Ophthalmol* 2019; 201:54–62.

Godefrooij D, Gans R, Imhof S, Wisse R. Nationwide reduction in the number of corneal transplantations for keratoconus following the implementation of crosslinking. *Acta Ophthalmol.* (2016) 94:675–8. doi: 10.1111/aos.13095.

Gokul A, Patel D, McGhee CN. John Nottingham's 1854 landmark treatise on conical cornea considered in the context of the current knowledge of keratoconus. *Cornea.* (2016) 35:673–8. doi: 10.1097/ICO.0000000000000801.

Gomes JAP, Rodrigues PF, Lamazales LL (2022) Keratoconus epidemiology: a review. *Saudi J Ophthalmol* 36(1):3–6.

Gomes, J. A. P., Tan, D., Rapuano, C. J., Belin, M. W., Ambrósio, R., Guell, J. L., ... & Wilson, S. E. (2015). Global consensus on keratoconus and ectatic diseases. *Cornea*, 34(4), 359–369.

Gondhowiardjo TD, Van Haeringen NJ. Corneal aldehyde dehydrogenase, glutathione reductase, and glutathione s- transferase in pathologic corneas. *Cornea* 1993; 12:310–4.

- Gonzalez V, McDonnell PJ. Computer-assisted corneal topography in parents of patients with keratoconus. *Arch Ophthalmol* 1992; 110:1412-4.
- Gordon-Shaag A, Millodot M, Essa M, Garth J, Ghara M, Shneur E. Is consanguinity a risk factor for keratoconus? *Optom Vis Sci* 2013; 90:448-54.
- Gore DM, Leucci MT, Anand V, Fernandez-Vega Cueto L, Arba Mosquera S, Allan BD. Combined wavefront-guided transepithelial photorefractive keratectomy and corneal crosslinking for visual rehabilitation in moderate keratoconus. *J Cataract Refract Surg* 2018; 44:571-80.
- Gorshkova E, Sevost'ianov E. Epidemiology of keratoconus in the Urals. *Vestn Oftalmol.* (1998) 114:38- 40.
- Grzybowski A, McGhee CNJ. The early history of keratoconus prior to Nottingham's landmark 1854 treatise on conical cornea: a review. *Clin Exp Optom* 2013. 96:140-5. <https://doi.org/10.1111/cxo.12035>.
- Hashemi H et al. *Cornea*. 2020;39(2):263-270.
- Hashemi H, Heydarian S, Hooshmand E, Saatchi M, Yekta A, Aghamirsalim M, et al. The Prevalence and risk factors for keratoconus: a systematic review and meta-analysis. *Cornea*. (2020) 39:263-70.
- Hashemi H, Khabazkhoob M, Fotouhi A. Topographic keratoconus is not rare in an Iranian population: the Tehran Eye Study. *Ophthalmic Epidemiol.* (2013) 20:385-91.
- Hersh, P. S., Greenstein, S. A., & Fry, K. L. (2011). Corneal collagen crosslinking for keratoconus and corneal ectasia: One-year results. *Journal of Cataract & Refractive Surgery*, 37(1), 149-160.
- Jonas JB, Nangia V, Matin A, Kulkarni M, Bhojwani K (2009) Prevalence and associations of keratoconus in rural maharashtra in central India: the central India eye and medical study. *Am J Ophthalmol* 148(5):760-765.
- Kandel H, Pesudovs K, Watson SL. Measurement of quality of life in keratoconus. *Cornea* 2020; 39:386-93.
- Kanski, J. J., & Bowling, B. (2015). *Kanski's Clinical Ophthalmology: A Systematic Approach*. 8th Edition. Elsevier.
- Kasparova EA. Pathogenetic basis for treatment of primary keratoconus by a combined method of excimer laser surgery (combination of photorefractive and phototherapeutic keratectomy). *Vestn Oftalmol* 2002; 118:21-215.
- Kemp EG, Lewis CJ. Immunoglobulin patterns in keratoconus with particular reference to total and specific IgE levels. *Br J Ophthalmol* 1982; 66:717-20.
- Kennedy RH, Bourne WM, Dyer JA: A 48-year clinical and epidemiologic study of keratoconus. *Am J Ophthalmol*. 2002, 101:267-73.
- Kiliç R, Cumurcu T, Sancaktar E, Evliyaoğlu O, Sezer H. Systemic prolidase activity and oxidative stress in keratoconus. *Curr Eye Res* 2016; 41:28-33.
- Kim KH, Mian SI. Refractive approaches to visual rehabilitation in patients with keratoconus. *Curr Opin Ophthalmol* 2020; 31:261-7.
- Klyce, S. D. (2009). Chasing the suspect: Keratoconus. *British Journal of Ophthalmology*, 93(7), 845-847.
- Konda S, Ambati BK. Intracorneal ring segments followed by toric pseudoaccommodating IOL for treatment of patients with corneal ectasia and cataract. *Am J Ophthalmol Case Reports* 2020;18.
- Koppen C, Kreps EO, Anthonissen L, Van Hoey M, Dhuhghaill SN, Vermeulen L. Scleral lenses reduce the need for corneal transplants in severe keratoconus. *Am J Ophthalmol* 2018; 185:43-7.
- Lapeyre G, Fournie P, Vernet R, Roseng S, Malecaze F, Bouzigon E, et al. Keratoconus

prevalence in families: a French study. *Cornea* 2020; 39:1473–9.

Leal F, Lipener C, Chalita MR, Uras R, Campos M, H'ofling-Lima AL. Lente de contato de material híbrido em pacientes com ceratocone e astigmatismo mi'opico composto. *Arq Bras Oftalmol* 2007; 70:247–54.

Lee A, Sakhalkar MV. Ocular manifestations of Noonan syndrome in twin siblings: a case report of keratoconus with acute corneal hydrops. *Indian J Ophthalmol* 201- 62:1171. 3.

Lim L, Lim EWL. Current perspectives in the management of keratoconus with contact lenses. *Eye* 2020; 34:2175–96.

Lim N, Vogt U. Characteristics and functional outcomes of 130 patients with keratoconus attending a specialist contact lens clinic. *Eye* 2002; 16:54–9.

Ling JJ, Mian SI, Stein JD, Rahman M, Poliskey J, Woodward MA. Impact of scleral contact lens use on the rate of corneal transplantation for keratoconus. *Cornea* 2021; 40:39–42.

Martinez-Abad A, Piñero DP. *J Cataract Refract Surg*. 2017;43(9):1213-1227.

Mathan JJ, Gokul A, Simkin SK, Meyer JJ, Patel DV, McGhee CNJ. Topographic screening reveals keratoconus to be extremely common in Down syndrome. *Clin Exp Ophthalmol* 2020; 48:1160–7.

Maya Tong C, Van Dijk K, Melles GRJ. Update on Bowman layer transplantation. *Curr Opin Ophthalmol* 2019; 30:249–55.

McMahon TT, Szczotka-Flynn L, Barr JT, Anderson RJ, Slaughter ME, Lass JH, et al. A new method for grading the severity of keratoconus: The Keratoconus Severity Score (KSS). *Cornea* 2006; 25:794–800.

McMonnies, C. W. (2015). Abnormal rubbing and, keratoconus. *Cornea* 1001-GG7, (12)34.1001-GG7, (12)34.

Mounir A, Mostafa EM. Combined accelerated corneal collagen crosslinking and intrastromal Kerarings implantation for treatment of advanced superior keratoconus. *GMS Ophthalmol Cases* 2020;10.

Navel V, Malecaze J, Pereira B, Baker JS, Malecaze F, Sapin V, et al. Oxidative and antioxidative stress markers in keratoconus: a systematic review and meta- analysis. *Acta Ophthalmol* 2020;99: e777–94.

O'Brart DPS. Corneal collagen crosslinking for corneal ectasias: A review. *Eur J Ophthalmol* 2017; 27:253–69. <https://doi.org/10.5301/ejo.5000916>.

Özalp O, Atalay E, Yıldırım N (2021) Prevalence and risk factors for keratoconus in a university-based population in Turkey. *J Cataract Refract Surg* 47(12):1524–1529.

Papali'i-Curtin A, Cox R, Ma T, Woods L, Covello A, Hall R. Keratoconus prevalence among high school students in New Zealand. *Cornea*. (2019) 38:1382–9.doi: 10.1097/ICO.0000000000002054.

Park C, Lee J, Gore P, Lim C, Chuck R. Keratoplasty in the United States. *Ophthalmology*. (2015) 122:2432–42.

Peña-García P, Sanz-Díez P, Durán-García ML. Keratoconus management guidelines. *Int J Keratoconus Ectatic Corneal Dis* 2015; 4:1–39.

Perry HD, Buxton JN, Fine BS. Round and oval cones in keratoconus. *Ophthalmology* 1980; 87:905–9.

Pilskalns B, Fink BA, Hill RM. Oxygen demands with hybrid contact lenses. *Optom Vis Sci* 2007; 84:334–42.

Rabinowitz YS, Galvis V, Tello A, Rueda D, García JD. Genetics vs chronic corneal mechanical trauma in the etiology of keratoconus. *Exp Eye Res* 2021; 202:108328.

Rabinowitz, Y. S. (1GG8). Keratoconus. Survey of, *Ophthalmology*, 42(4).31G-2G7.

Robertson I. Keratoconus and the Ehlers Danlos syndrome: a new aspect of keratoconus. *Med J Aust* 1975; 1:571–3.

Romero-Jiménez M, Santodomingo-Rubido J, Flores- Rodríguez P, González- M'eijome JM. Short-term corneal changes with gas-permeable contact lens wear in keratoconus subjects: a comparison of two fitting approaches. *J Optom* 2015;8: 48–55.

Romero-Jiménez M, Santodomingo-Rubido J, González- M'eijome JM. An assessment of the optimal lens fit rate in keratoconus subjects using three-point- touch and apical touch fitting approaches with the rose K2 lens. *Eye Contact Lens* 2013; 39:269–72.

Rosenthal P, Croteau A. Fluid-ventilated, gas-permeable scleral contact lens is an effective option for managing severe ocular surface disease and many corneal disorders that would otherwise require penetrating keratoplasty. *Eye Contact Lens* 2005; 31:130–4.

Rubinstein MP, Sud S. The use of hybrid lenses in management of the irregular cornea. *Cont Lens Anterior Eye* 1999; 22:87–90.

Salman A, Darwish T, Ghabra M, Kailani O, Haddeh Y, Askar M et al (2022) Prevalence of keratoconus in apopulation-based study in Syria. *J Ophthalmol* 2022:6064533

Sandvik G, Thorsrud A, Råen M, Østern A, Sæthre M, Drolsum L. Does corneal collagen cross-linking reduce the need for keratoplasties in patients with keratoconus? *Cornea*. (2015) 34:991–5.

Santodomingo-Rubido J, Carracedo G, Suzaki A, Villa- Collar C, Vincent SJ, Wolffsohn JS (2022) Keratoconus: an updated review. *Cont Lens Anterior Eye* 45(3):101559

Saraç "O, Kars ME, Temel B, Çağıl N. Clinical evaluation of different types of contact lenses in keratoconus management. *Contact Lens Anterior Eye* 2019;42: 482–6.

Segal O, Barkana Y, Hourovitz D, Behrman S, Kamun Y, Avni I, et al. Scleral contact lenses

may help where other modalities fail. *Cornea* 2003; 22:308–10.

Shetty R, Sharma A, Pahuja N, Chevour P, Padmajan N, Dhamodaran K, et al. Oxidative stress induces dysregulated autophagy in corneal epithelium of keratoconus patients. *PLoS ONE* 2017;12: e0184628.

Sinha A, Goel S, Gupta V, Kumawat D, Sahay P. iTrace – A Ray Tracing Aberrometer. *Delhi J Ophthalmol* 2019;30:72-75.

Sorkin N, Varssano D. Corneal collagen crosslinking: A systematic review. *Ophthalmologica* 2014; 232:10–27.

Srivastava OP, Chandrasekaran D, Pfister RR. Molecular changes in selected epithelial proteins in human keratoconus corneas compared to normal corneas. *Mol Vis* 2006; 12:1615–25.

Sultan P, Dogan C, Iskeleli G. A retrospective analysis of vision correction and safety in keratoconus patients wearing Toris K soft contact lenses. *Int Ophthalmol* 2016; 36:799–805.

Sykakis, E., Karim, R., Evans, J. R., Bunce, C., Amissah- &, Arthur, K. N. Patwary, S. (2015). Corneal collagen cross-linking for treating. keratoconus. *Cochrane Database of Systematic Reviews*.

Ting D, Sau C, Srinivasan S, Ramaesh K, Mantry S, Roberts F. Changing trends in keratoplasty in the West of Scotland: a 10-year review. *Br J Ophthalmol*. (2012) 96:405–8. doi: 10.1136/bjophthalmol-2011-300244.

Torres Netto E, Al-Otaibi W, Hafezi N, Kling S, Al-Farhan H, Randleman J, et al. Prevalence of keratoconus in paediatric patients in Riyadh, Saudi Arabia. *Br J Ophthalmol*. (2018) 102:1436–41.

Torres Netto EA, Al-Otaibi WM, Hafezi NL, Kling S, Al- Farhan HM, Randleman JB et al (2018) Prevalence of keratoconus in paediatric patients in Riyadh. *Saudi Arabia Br J Ophthalmol* 102(10):1436–1441.

Tuft SJ, Hassan H, George S, Frazer DG, Willoughby CE, Liskova P. Keratoconus in 18 pairs of twins. *Acta Ophthalmol* 2012;90: e482–6.

Vega Estrada A, Sanz Díez P, Ali'ó JL. Keratoconus grading and its therapeutic implications. In: Ali'ó J. (eds) *Keratoconus. Essentials in ophthalmology*, 2017, p. 177–84 . doi: 10.1007/978-3-319-43881-8_15.

Wang Y, Rabinowitz YS, Rotter JI, Yang H. Genetic epidemiological study of keratoconus: Evidence for major gene determination. *Am J Med Genet* 2000; 93:403.

Wollensak, G., Spoerl, E., & Seiler, T. (2003). Riboflavin/ultraviolet-A-induced collagen crosslinking for the treatment of keratoconus. *American Journal of Ophthalmology*, 135(5), 620–627.

Wyman A. Benedict Duddell: pioneer oculist of the 18th century. *J R Soc Med.* (1992) 85:412–5. Xiong S, Lv M, Zou H, Zhu J, Lu L, Zhang B, Deng J, Yao C, He X, Xu X. Comparison of Refractive Measures of Three Autorefractors in Children and Adolescents. *Optom Vis Sci.* 2017 Sep;94(9):894-902.

Yam GHF, Fuest M, Zhou L, Liu YC, Deng L, Chan ASY, *et al.* Differential epithelial and stromal protein profiles in cone and non-cone regions of keratoconus corneas. *Sci Rep* 2019; 9:2965.

Yilmaz I, Ozcelik F, Basarir B, Demir G, Durusoy G, Taskapili M. Clinical performance of toris K contact lens in patients with moderate to advanced keratoconus: A real- life retrospective analysis. *J Ophthalmol* 2016;2358901.

Zadnik K, Barr JT, Edrington TB, Everett DF, Jameson M, McMahon TT, *et al.* Baseline findings in the collaborative longitudinal evaluation of keratoconus (CLEK) study. *Invest Ophthalmol Vis Sci* 1998; 39:2537–46.